

## Hydro geochemical Insights and Groundwater Quality Assessment of Pulivendula Mandal, Kadapa, Andhra Pradesh

C. Krupavathi, S. Srinivasa Gowd\*, P. Ravi Kumar, G.Nagalakshmi Devi, U. Lahari

Department of Geology, Yogi Vemana University, Kadapa, Andhra Pradesh, India

### ABSTRACT

Several hydro geochemical processes controlling the geochemistry of aquifers, the suitability of groundwater for agriculture, and routine use were assessed in the Pulivendula area during the current survey. To characterize the main physicochemical components of the aquifer water, a total of 24 drilling locations were selected. The findings showed that the groundwater had an alkaline pH. Due to the ideal limits of total dissolved solids, magnesium ( $Mg^{2+}$ ), sodium ( $Na^+$ ), potassium ( $K^+$ ), sulphate ( $SO_4^{2-}$ ), and chloride ( $Cl^-$ ), the majority of samples fall within the appropriate range. Residential use of groundwater should be avoided in areas when EC, total hardness, calcium ( $Ca^{2+}$ ), and nitrate ( $NO_3^-$ ) slightly exceed recommended standards. With very few exceptions, the groundwater in the research region is appropriate for human consumption, according to the analytical findings. The Piper diagram was utilized to analyze all samples to identify the primary hydro geochemical components found in groundwater. In this area, cations predominate in groundwater with  $Ca^{2+} > Na^+ > Mg^{2+} > K^+$ , followed by anions  $HCO_3^- > Cl^- > SO_4^{2-} > F^-$ . According to the Gibbs plot study, the primary hydro geochemical mechanism controlling the geochemistry of the groundwater in this area is the rock-aquifer interaction. It was discovered that the greatest number of water samples was ideal for cultivation.

**Key words:** Hydrogeochemistry, Groundwater Quality Assessment, Pulivendula, YSR District

### 1. INTRODUCTION

The main supply of water for agriculture, industry, home usage, and drinking is groundwater [1-4]. The increasing demand for groundwater may be attributed to several factors, including the rapid rise of the population, the development of irrigation, and the increasing trend of industrialization [5-8]. The main causes of anthropogenic contamination, which comes from sources like industrial effluents, agricultural fertilizers, municipal waste water, septic tank effluent, landfills, and animal wastes, are also responsible for the degradation of groundwater quality. A high level of human health and agricultural productivity are directly correlated with water quality [9-11]. In many regions of India, the lack of clean, drinkable water has become one of the most important developmental challenges in recent years [12-14]. Both natural processes and human activity have an impact on groundwater quality, which is a global environmental concern [15,16].

Human health is greatly impacted by the quality of groundwater. A little shift in the quality of the water indicates that the water ecosystem is not working properly [17,18]. Because of weathering from source rocks, human activity, and regional environmental and ecological factors, physical and chemical soluble characteristics have a significant impact on groundwater quality [19-21]. Excessive accumulations of pollutants on the land and contamination of available surface and subsurface water resources were caused by the careless disposal of anthropogenic, agricultural, and mining wastes, the unplanned application of agrochemicals and fertilizers, and public ignorance of environmental concerns [22]. Numerous factors influence the chemical makeup of groundwater, such as precipitation composition, the geology and mineralogy of aquifers and watersheds, and the geological processes that occur inside aquifers [23,24].

Different kinds of water result from the interplay of all these variables. A sustainable development of water resources requires an understanding of the geochemical evolution of groundwater [25]. Geological formations and human activity have a significant impact on the physical and chemical characteristics that determine how the quality of groundwater varies in a given location [26,27]. Water of poor quality has a negative impact on human health and plant development. Soil, lithology, and geology are some of the key elements that affect water quality. Depending on variations in geological formations, water quality can change [28]. Due to water contamination brought on by the disposal of overburden in and near mining zones [29,30]. The depth of the water table, location, and season all affect how pure the groundwater is [31]. The amount and make-up of dissolved solids in water determine the quality in the main [32]. The weathering of rocks and subsequent contributions from soils can both contribute to the area's groundwater quality degradation [33,34]. To determine if the groundwater in and around Pulivendula, Kadapa district, Andhra Pradesh, is suitable for human consumption, this article examines a number of chemical parameters.

### \*Corresponding author:

S. Srinivasa Gowd

E-mail: ssgowd@gmail.com

ISSN NO: 2320-0898 (p); 2320-0928 (e)

DOI: 10.22607/IJACS.2025.1301004

Received: 09<sup>th</sup> November 2024;

Revised: 28<sup>th</sup> December 2024;

Accepted: 02<sup>th</sup> January 2025;

Published: 02<sup>th</sup> February 2025



## 2. METHODOLOGY

To evaluate the groundwater quality, 24 groundwater samples were taken in the Pulivendula region from chosen bore wells (deep aquifer) and dug wells (shallow aquifer) during the pre-monsoon and post-monsoon seasons of 2024 (May and October). These 1-liter plastic containers that had been previously cleaned were used to gather the water samples. Groundwater sample physicochemical analyses were conducted using American Public Health Association (APHA) standard operating procedures. Following the collection of groundwater samples in the field, the physical characteristics, such as pH, electrical conductivity (EC), and total dissolved solids (TDS), were determined right away. The spectrophotometer was used to analyze the sulfate ( $\text{SO}_4^{2-}$ ), fluoride ( $\text{F}^-$ ), and nitrate ( $\text{NO}_3^-$ ), while the titrimetric method was used to analyze the bicarbonates ( $\text{HCO}_3^-$ ), calcium ( $\text{Ca}^{2+}$ ), magnesium ( $\text{Mg}^{2+}$ ), and chloride ( $\text{Cl}^-$ ). The flame photometer was used to estimate the chemical parameters such as sodium ( $\text{Na}^+$ ) and potassium ( $\text{K}^+$ ). Tables 1 and 2 show the physical characteristics and results of the chemical analysis of groundwater samples.

### 2.1. Location Accessibility

The area lies between Pulivendula situated between parallels of  $78^\circ 08' 0''$  to  $78^\circ 19' 0''$  E L and  $14^\circ 13' 30''$  to  $14^\circ 30' 0''$  N L with intended boundary falling in Survey of India topographic sheet 57J 03 on 1:50,000 scale covering an of over  $720 \text{ km}^2$ . The study area is shown in Figure 1.

### 2.2. Physiographic and Climate

The research area's southern and central regions are made up of lowlands, ridges, and tall hills. In the research region, contour values vary from 200 to 600 m. There is a  $1-35^\circ$  range in the slope category. The Mogamureru River is the sole seasonal river in the research region, receiving its water during the monsoon season (June to October) and flowing northwest through the southern portion of the study area. The same region is also covered by a few large tanks. The average annual temperature is  $32^\circ\text{C}$ , with 100–150 cm of precipitation and a dry environment. December is thought to be the coldest month ( $25^\circ\text{C}$ ), and May is thought to be the warmest ( $45^\circ\text{C}$ ). Every crop is longer than 120 days.

### 2.3. Study Area

In the drought-stricken and economically challenged Kadapa district of Andhra Pradesh, Pulivendula Mandal is an overfished Mandal. Anantapur district borders the Mandal on the south, Vemula Mandal on the east, Thondur Mandal on the north, and Anantapur district on the west. It is situated in the Kadapa district's western region [Figure 1]. The research region, which covered  $182.6 \text{ km}^2$  geographically, had a dendritic structure as a result of geological formations.

### 2.4. Geology

The research region is a part of Southern India's most important Purana basins. The Lower Cuddapah Super group's Papaghni and

**Table 1:** Physicochemical parameters concentrations of the study area during pre-monsoon

Sample No.	Well depth (M)	pH	TDS	EC	TH	$\text{Ca}^{2+}$	$\text{Mg}^{2+}$	$\text{K}^+$	$\text{Na}^+$	$\text{Cl}^-$	$\text{NO}_3^-$	$\text{SO}_4^{2-}$	$\text{HCO}_3^-$	$\text{F}^-$
1	20	7.9	1208.2	724.9	732.5	137.7	94.4	2.84	24.6	163.5	32.9	265.4	453.6	1.7
2	25	8.4	1219.6	731.7	624.6	95.3	93.9	1.96	37.6	175.4	51.4	265.4	460.8	1.9
3	35	8.2	1286.5	771.9	680.2	113.1	96.6	0.94	27.6	175.0	50.4	233.4	548.3	2.5
4	20	8	1090.0	654.0	633.2	122.7	79.4	1.62	9.6	148.5	17.9	250.4	438.6	3.5
5	31	7.9	1330.8	798.5	831.7	152.7	109.4	2.8	39.6	178.5	47.9	280.4	468.6	4.3
6	35	7.65	1341.8	805.1	723.8	110.3	108.9	1.52	52.6	190.4	66.4	280.4	475.8	4.1
7	39	8.15	1408.3	845.0	779.4	128.1	111.6	1.62	42.6	190.0	65.4	248.4	563.3	4.4
8	20	7.68	1105.5	663.3	611.4	105.7	84.4	1.18	14.6	153.5	22.9	255.4	443.6	1.0
9	27	8.39	1116.7	670.0	503.5	63.3	83.9	2.78	27.6	165.4	41.4	255.4	450.8	1.0
10	55	7.87	1184.9	711.0	559.1	81.1	86.6	2.54	17.6	165.0	40.4	223.4	538.3	3.0
11	34	7.95	1065.3	639.2	512.2	90.7	69.4	1.62	46.9	138.5	37.9	240.4	428.6	3.5
12	60	7.47	1293.6	776.2	799.3	172.7	89.4	1.21	69.6	149.5	56.4	214.0	528.6	4.0
13	45	7.39	1210.5	726.3	725.1	160.3	78.9	2.98	72.6	100.4	55.4	140.4	585.8	3.9
14	38	7.99	1205.6	723.4	755.8	168.1	81.6	1.78	72.6	90.0	22.9	128.4	623.3	4.1
15	30	7.75	1158.6	695.2	627.3	146.7	63.4	2.12	43.6	123.5	52.9	188.0	502.6	3.5
16	26	8.31	1095.7	657.4	553.1	134.3	52.9	2.24	46.6	74.4	71.4	114.4	559.8	4.0
17	42	8.43	1112.6	667.5	583.8	142.1	55.6	1.36	36.9	64.0	70.4	102.4	597.3	4.3
18	30	8.2	1088.1	652.8	595.8	140.7	59.4	1.54	59.6	128.5	27.9	230.4	418.6	3.6
19	34	7.05	1322.0	793.2	647.1	128.3	79.4	2.38	62.6	183.5	46.4	271.4	528.3	2.8
20	28	8.04	1192.0	715.2	623.6	136.1	68.9	2.2	62.6	195.4	45.4	239.4	418.6	2.7
21	40	8.1	1260.4	756.2	581.2	114.7	71.6	0.92	33.6	195.0	42.9	256.4	518.6	2.9
22	45	8.5	1321.1	792.6	475.1	102.3	53.4	1.02	36.6	158.5	61.4	286.4	575.8	2.3
23	38	8.07	1361.0	816.6	451.6	110.1	42.9	2.18	26.9	170.4	60.4	286.4	613.3	2.8
24	37	8.4	1201.6	720.9	459.2	108.7	45.6	1.94	49.6	170.0	27.9	254.4	492.6	3.1

EC: Electrical conductivity, TH: Total hardness, TDS: Total dissolved solids,  $\text{SO}_4^{2-}$ : Sulphate,  $\text{F}^-$ : Fluoride,  $\text{NO}_3^-$ : Nitrate,  $\text{HCO}_3^-$ : Bicarbonates,  $\text{Ca}^{2+}$ : Calcium,  $\text{Mg}^{2+}$ : Magnesium,  $\text{Cl}^-$ : Chloride,  $\text{Na}^+$ : Sodium,  $\text{K}^+$ : Potassium

**Table 2:** Physicochemical parameters concentrations of the study area during post-monsoon

Sample No.	Well depth (M)	pH	TDS	EC	TH	Ca <sup>2+</sup>	Mg <sup>2+</sup>	K <sup>+</sup>	Na <sup>+</sup>	Cl <sup>-</sup>	NO <sub>3</sub> <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>	HCO <sub>3</sub> <sup>-</sup>	F <sup>-</sup>
1	25	7.4	1060.3	636.2	600.2	117.7	74.4	3.24	24.6	143.5	12.9	245.4	433.6	1.7
2	30	7.9	1070.8	642.5	492.3	75.3	73.9	2.36	17.6	155.4	31.4	245.4	440.8	1.1
3	40	7.7	1139.0	683.4	547.9	93.1	76.6	1.34	7.6	155.0	30.4	213.4	528.3	3.0
4	25	7.5	942.1	565.3	500.9	102.7	59.4	2.02	35.0	128.5	32.5	230.4	418.6	3.5
5	36	7.4	1182.5	709.5	699.4	132.7	89.4	3.2	19.6	158.5	27.9	260.4	448.6	3.9
6	40	7.15	1191.2	714.7	591.5	90.3	88.9	1.92	32.6	170.4	46.4	260.4	455.8	4.9
7	44	7.65	1256.5	753.9	647.1	108.1	91.6	2.02	22.6	170.0	45.4	228.4	543.3	2.2
8	25	7.18	990.6	594.4	479.1	85.7	64.4	1.58	35.8	133.5	2.9	235.4	423.6	3.7
9	32	7.89	968.1	580.9	371.2	43.3	63.9	3.18	7.6	145.4	21.4	235.4	430.8	3.8
10	60	7.37	1033.5	620.1	426.8	61.1	66.6	2.94	45.8	145.0	20.4	203.4	518.3	1.2
11	39	7.45	936.9	562.1	379.9	70.7	49.4	2.02	26.9	118.5	17.9	220.4	408.6	2.6
12	55	8.4	1161.9	697.2	667.0	152.7	69.4	1.61	49.6	129.5	36.4	194.0	508.6	2.7
13	50	8.3	1080.3	648.2	592.8	140.3	58.9	3.38	52.6	80.4	35.4	120.4	565.8	4.1
14	43	7.49	1072.7	643.6	623.5	148.1	61.6	2.18	52.6	70.0	2.9	108.4	603.3	1.6
15	35	7.25	1039.4	623.6	495.0	126.7	43.4	2.52	23.6	103.5	32.9	168.0	482.6	1.7
16	31	7.81	940.7	564.4	420.8	114.3	32.9	2.64	26.6	54.4	51.4	94.4	539.8	2.5
17	47	7.93	960.5	576.3	451.5	122.1	35.6	1.76	16.9	44.0	50.4	82.4	577.3	4.0
18	35	7.7	958.9	575.3	463.5	120.7	39.4	1.94	39.6	108.5	7.9	210.4	398.6	4.1
19	39	7.69	1173.6	704.1	514.8	108.3	59.4	2.78	42.6	163.5	26.4	251.4	508.3	4.4
20	33	7.54	1068.7	641.2	491.3	116.1	48.9	2.6	42.6	175.4	25.4	219.4	398.6	2.9
21	45	7.6	1138.7	683.2	448.9	94.7	51.6	1.32	13.6	175.0	22.9	236.4	498.6	3.0
22	50	8	1182.9	709.8	342.8	82.3	33.4	1.42	16.6	138.5	41.4	266.4	555.8	3.9
23	43	7.57	1190.1	714.1	319.3	90.1	22.9	2.58	6.9	150.4	40.4	266.4	593.3	5.4
24	42	7.9	1032.1	619.3	538	88.7	25.6	2.34	29.6	150.0	7.9	234.4	472.6	5.5

EC: Electrical conductivity, TH: Total hardness, TDS: Total dissolved solids, SO<sub>4</sub><sup>2-</sup>: Sulphate, F<sup>-</sup>: Fluoride, NO<sub>3</sub><sup>-</sup>: Nitrate, HCO<sub>3</sub><sup>-</sup>: Bicarbonates, Ca<sup>2+</sup>: Calcium, Mg<sup>2+</sup>: Magnesium, Cl<sup>-</sup>: Chloride, Na<sup>+</sup>: Sodium, K<sup>+</sup>: Potassium

Chitravati groups make up the stratigraphically studied region [35]. Lithologically, this area is made up of enormous limestone, dolomite, basalt, shale, conglomerate, and some of the most common intrusions (sills) [Figures 2 and 3]. The pediment, pediplain, denudational hills, and structural hills are the main geomorphic characteristics in this area. Groundwater is present in the worn areas of the Chitravathi and Papagani rock formations. The existence of multiple joints, fractures, and fissure zones in this kind of rock is primarily responsible for the water found in drilled wells in this area. There are good aquifers made of massive limestone, shale, and quartzite, and permanent groundwater levels are often shallow. The depth of the water varies from 10 m to 50 m [36]. Important crops include sweet lime and paddy; banana plantations also yield seasonal fruits like lemon and sweet lime. There is an average of 600–650 mm of rainfall per year, and the north-east monsoon and south-west wind both have an impact.

### 2.5. Geographical Information System (GIS) Analysis

Understanding water quality study and map development requires the use of GIS. The current investigation was conducted using Arc GIS 10.3 and topographical sheets. Examine the area map created using 1:50,000 scales then create base maps using the World Geodetic System (WGS) 1984 coordinate system. Throughout the study region, the

sample collection mapping was delineated using GPS measurements. Using bar graphs and charts, the extended module of Arc GIS 10.2 was utilized to create a range of distribution maps on hardness, pH, and other important cations and anions.

## 3. MECHANISM CONTROL FACTORS

### 3.1. Gibbs Diagram and Scatter Plots

Evaporation is a crucial factor that can change groundwater chemistry in addition to precipitation and the water-rock interaction (rock dominance) [37]. Because the majority of the samples occur within the carbonate weathering zone, the scatter plot between the Ca<sup>2+</sup>/Na<sup>+</sup> and Mg<sup>2+</sup>/Na<sup>+</sup> values indicates that rock–water interaction is often related with silicate weathering [38].

### 3.2. Hydro-facial

The chemical makeup of groundwater solutions observed in hydrogeochemical systems is referred to as hydrogeochemical facies [39]. Determining the facies is helpful in comprehending the connections and similarities between various ions found in the groundwater of an aquifer as well as the impact of chemical interactions between minerals and groundwater that take place within a lithological framework. Although there are several methods for




Figure 1: Study area.



Figure 2: Stramatolites in Vempalle formation.



Figure 3: Quartzite hill and Vempalle formations just below quartzite hills and shale beds.

interpreting groundwater hydrogeochemistry, including graphical and statistical analysis, Piper and Gibbs diagrams are frequently utilized to determine hydrogeochemical facies [40]. Six fields [41] are used to depict this diagram:  $\text{Ca}^{2+}$ -  $\text{HCO}_3^-$  type,  $\text{Na}^+$ -  $\text{Cl}^-$  type,  $\text{Ca}^{2+}$ -  $\text{Mg}^{2+}$ -  $\text{Cl}^-$  type,  $\text{Ca}^{2+}$ -  $\text{Na}^+$ -  $\text{HCO}_3^-$  type,  $\text{Ca}^{2+}$ -  $\text{Cl}^-$  type, and  $\text{Na}^+$ -  $\text{HCO}_3^-$  type.

### 3.3. Saturation Index (SI)

The SI studies the chemical equilibrium of processes involving minerals and aqueous species [42]. The groundwater sample saturation indices were computed using Phreeqc Interactive version 3.4. The formula that defines the SI system is  $\text{SI} = \log_{10} \text{IAP/K}$  mineral, where IAP stands for the ion activity product and K mineral for the solubility constant of the mineral. The soluble mineral in the water sample is fully saturated when SI is equal to

0 [43-45]. Positive specific gravity (SI) values suggest that the sample is super-saturated, which will lead to the mineral tending to precipitate; negative SI values indicate that the sample is under-saturated, and the mineral will dissolve.

## 4. RESULTS AND DISCUSSION

Chemistry of groundwater Evocative data are calculated and compared with the World Health Organization [46] drinking water quality guidelines [Tables 1 and 2] to assess whether groundwater is suitable for human use. The pH scale runs from 7.0 to 8.5, 7.15 to 8.4, and the mean value of 8 indicates that the groundwater is alkaline. The groundwater samples have EC values ranging from 639.2 to 845  $\mu\text{S}/\text{cm}$  and 562.1 to 753.9  $\mu\text{S}/\text{cm}$ , respectively, with a

mean of 729.5  $\mu\text{S}/\text{cm}$  for both seasons. A higher EC value denotes a greater salinity and mineral content in groundwater that has low runoff, high infiltration, and a specific type of discharge [47]. On the other hand, high elevated terrain, heavy runoff, low infiltration and recharge water type, and low salt enrichment are typically linked to low EC values. All inorganic salts that indicate the salinity of the water and its acceptability for human consumption are included in the total dissolved solids (TDS) in water [37]. The TDS in groundwater samples has a mean value of 1215.9 mg/L for both seasons and varies from 1065.3 to 1408.3 mg/L and 936.9 to 1256.5 mg/L [Table 3]. Because nearly all of the samples are beyond the groundwater's EC and fall below the permissible limit of 500 mg/L, they cannot be used for irrigation or drinking. Groundwater water's total hardness (TH) is a result of the presence of metal ions such as  $\text{Ca}^{2+}$  and magnesium. Higher quantities of water containing TH (>300 mg/L) have been linked to health concerns, including renal difficulties. Hard water is therefore inappropriate for use in homes. Consequently, it is discovered that groundwater is unsuitable for agriculture, irrigation, and drinking in the majority of the research region [48].

#### 4.1. Cation Chemistry

Because it promotes the formation of teeth and bones, the concentration of  $\text{Ca}^{2+}$  cation in groundwater is a crucial aspect of groundwater chemistry. Table 2 indicates that the  $\text{Ca}^{2+}$  content in the analyzed samples fluctuates between 63.3 and 172.7 mg/L and between 63.3 and 172.7 mg/L [Figure 4]. The average values for both seasons are 123.6 and 123.6 mg/L, respectively. When compared to the World Health Organization (WHO) guidelines, it shows a slightly higher  $\text{Ca}^{2+}$  concentration.  $\text{Ca}^{2+}$ -enriched groundwater is typically found in rocks and minerals such as shale, dolomite, and limestone, as well as in minerals like pyroxene, amphibole, and plagioclase. Moreover, ion exchange and the existence of carbon dioxide in the soil zone are additional sources of  $\text{Ca}^{2+}$  entering the groundwater [49]. In the same way, the majority of these rocks and minerals in the research region provide  $\text{Ca}^{2+}$  to the groundwater.

Magnesium is typically found in natural waterways alongside  $\text{Ca}^{2+}$ . It can also come from human (such as mining operations and industrial waste) or geogenic (seawater, ferro-magnesium rocks, and ion exchange) sources. According to Table 3, its content in the study region ranges from 42.9 to 111.6 mg/L and from 22.9 to 91.6 mg/L, with averages of 77.6 and 57.6 mg/L for each season. The majority of samples have  $\text{K}^+$  concentrations that are found to be below the allowable limits, and only a small number of samples have  $\text{K}^+$  concentrations that are found to be beyond the allowable limits as per the WHO guidelines [Table 2]. In the research region, cations were found to be more abundant in the following order:  $\text{Ca}^{2+} > \text{Na}^+ > \text{Mg}^{2+} > \text{K}^+$ .

#### 4.2. Anion Chemistry

The most significant chemical present in natural water is  $\text{HCO}_3^-$ .  $\text{HCO}_3^-$ , which contributes to the alkaline quality of groundwater, is often produced by the weathering of silicate rocks, although it may also arise from primary carbonate and calcareous rocks [50]. Its content in the region is well under the allowable limit, ranging from 418.6 to 623.3 mg/L and 398.6 to 603.3 mg/L [Table 3], with averages of 509.7 and 489.7 mg/L for each season.

A major anion present in groundwater is  $\text{Cl}^-$ , which can come from a variety of sources, including human sources, leaching, and the weathering of various minerals [11]. Due to its high concentration in groundwater, it has a salty taste and can cause kidney stones, which is bad for human health [51]. The  $\text{Cl}^-$  content in the study region ranges from 64.0 to 195.4 mg/L and 44.0 to 175.4 mg/L [Table 3], with an average of 152.0 and 132 mg/L for each season. These results show that the groundwater in the area is safe to drink and is within allowable limits [46].

Geogenic and anthropogenic sources can produce  $\text{SO}_4^{2-}$ , which, at high concentrations, renders groundwater unfit for human consumption. Rich in gypsum material, carbonate sedimentary rocks are the geogenic source of  $\text{SO}_4^{2-}$ . With an average of 229.6 mg/L and 209.6 mg/L for both seasons, respectively, its concentration

**Table 3:** Descriptive statistics of the physicochemical parameters of groundwater in the Mogamureru River basin

Parameters	WHO international standard (WHO 2017)		Study area ranges (PRM)		Study area ranges (POM)	
	Desirable limits	Permissible limits	Min	Max	Min	Max
pH	6.5	8.5	7.05	8.5	7.15	8.4
EC	-	-	639.2	845	562.1	753.9
TDS	500	1500	1065.3	1408.3	936.9	1256.5
TH	100	500	451.6	831.7	319.3	699.4
$\text{HCO}_3^-$	-	-	418.6	623.3	398.6	603.3
$\text{SO}_4^{2-}$	200	400	102.4	286.4	82.4	266.4
$\text{Cl}^-$	200	600	64	195.4	44	175.4
$\text{NO}_3^-$	50	-	17.9	71.4	2.9	51.4
$\text{F}^-$	0.5	1.5	1	4.4	1.1	5.5
$\text{Ca}^{2+}$	75	100	63.3	172.7	63.3	172.7
$\text{Mg}^{2+}$	50	150	42.9	111.6	22.9	91.6
$\text{Na}^+$	-	200	9.6	72.6	6.9	52.6
$\text{K}^+$	-	12	0.92	2.98	1.32	3.38

WHO: World Health Organization, EC: Electrical conductivity, TH: Total hardness, TDS: Total dissolved solids,  $\text{SO}_4^{2-}$ : Sulphate,  $\text{F}^-$ : Fluoride,  $\text{NO}_3^-$ : Nitrate,  $\text{HCO}_3^-$ : Bicarbonates,  $\text{Ca}^{2+}$ : Calcium,  $\text{Mg}^{2+}$ : Magnesium,  $\text{Cl}^-$ : Chloride,  $\text{Na}^+$ : Sodium,  $\text{K}^+$ : Potassium, PRM: Pre monsoon, POM: Post monsoon

varies from 102 to 286.4 mg/L and 82.4 to 266.4 mg/L [Table 3]; this indicates that they are within the allowable limit in accordance with established guidelines [46; Table 2]. The dissolution of  $F^-$  in groundwater derived from  $F^-$ -bearing minerals is primarily responsible for the higher concentration of  $F^-$ . The dissolution of  $F^-$  in groundwater derived from  $F^-$ -bearing minerals is primarily responsible for the higher concentration of  $F^-$ . This process is influenced by a number of factors, including the source rocks, well depth, residential period, and favorable environments for the upward rise of deep-seated groundwater [52]. The groundwater samples exhibit a range of  $F^-$  concentrations, from 1 to 4.4 mg/L and 1.1 to 5.5 mg/L [Table 3], with averages of 3.1 mg/L and 3.2 mg/L for both seasons, respectively. This suggests that a majority of the study area's samples surpass the permission limit [46]. The geogenic processes of weathering, ion exchange, and leaching are most likely the cause of the samples with moderate-to-high  $F^-$  concentrations. Dental fluorosis is often brought on by ingesting groundwater that is high in  $F^-$  ( $>1.50$  mg/L) [53].

These days, concentrations of  $NO_3^-$ , particularly in places where industrialization, intensive agriculture, and population increase have been observed, are thought to be significant groundwater pollutants [54].  $NO_3^-$  can originate from either geological or human sources. The primary man-made sources of  $NO_3^-$  in groundwater include septic tanks, pipe leaks, and agricultural runoff resulting from the use of various fertilizers in human populated areas. Groundwater's high  $NO_3^-$  content results in health issues that

can progress to cancer, stomach issues, and Methaemoglobinemia (also known as "Blue baby" sickness) [55]. The study area's  $NO_3^-$  concentrations range from 17.9 to 71.4 mg/L and 2.9 to 51.4 mg/L, respectively [Table 3]; the average for both seasons is 46.5 mg/L, although groundwater samples slightly above the desired limit [46]. The statistical analysis with help of arc GIS of the research field is displayed in Figure 4.

The results show the dominance of anions as  $HCO_3^- > SO_4^{2-} > Cl^- > NO_3^- > F^-$  in the study area.

4.3. Gibbs Diagram and Scatter Plot

The bulk of the groundwater samples in our investigation, according to gibbs diagram (GD), was found in the rock-weathering dominance field [Figure 5] [37]. This implies that the primary natural factor influencing the chemistry of groundwater is the interaction between water and rock. The replenishment of fresh surface water into fresh groundwater is referred to as "rock-weathering dominance." Moreover, it implies that groundwater contains the bulk of dissolved carbonate minerals. Furthermore, it is evident that evaporation processes had an impact on groundwater, suggesting that there was a significant ion exchange between groundwater and solid components in surrounding sediment. Based on the analysis of the scatter plot research area, it was determined that the best governing mechanism for determining the hydrochemistry of the silicate-rock materials in the study region is silicate weathering [Figure 6].




Figure 4: Statistical analysis for pre and post-monsoon.




Figure 5: Gibb's diagrams for pre and post-monsoon.




Figure 6: Scatter plots for pre and post-monsoon.





Figure 7: Piper diagrams for pre and post-monsoon.



**Figure 8:** Saturation index.

#### 4.4. Hydro Geochemical Facies

An aquifer system's anions and cations' hydrogeochemical properties are shown in a piper diagram [41]. To determine the overall geochemical character and water quality, the concentrations of the anions ( $\text{Cl}^-$ ,  $\text{HCO}_3^-$ , and  $\text{SO}_4^{2-}$ ) and cations ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^+$ , and  $\text{K}^+$ ) were shown in this diagram. Plotted sample data in this study indicate that water samples fall into four different feldspars, with 90% of samples falling into the  $\text{Ca}^{2+}$ - $\text{Mg}^{2+}$ - $\text{HCO}_3^-$  facies and the remaining samples falling into other facies [Figure 7]. The  $\text{Ca}^{2+}$ - $\text{Mg}^{2+}$ - $\text{HCO}_3^-$  water type demonstrates that the predominant cations are  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ , whereas the major anion is  $\text{HCO}_3^-$ . The findings therefore show that weak acids ( $\text{HCO}_3^-$ ) and alkaline earth metals ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ) control the hydrochemistry of the groundwater.

#### 4.5. SI

Index of Saturation SI helps the groundwater system realize its reactive minerals since it is impacted by a variety of solutes that come from the contact with the adjacent sediments. According to SI standards, the majority of groundwater samples (5, 7, 10, 12, 13, 14, 15, 19, 20, 23, and 24) were supersaturated by dolomite mineral, whereas the rest of the samples are under saturated. Sample No. exhibits super saturation with calcite, whereas samples Nos. 5, 12, 13, 14, 15, 23, and 24 exhibit super saturation by dolomite mineral, and the other samples are under saturated [Figure 8].

### 5. CONCLUSIONS

The chemistry of groundwater is significantly influenced by a variety of processes, including evaporation throughout flow pathways, dissolution, and contact with rocks. Saturation indices for carbonate (calcite, dolomite, and aragonite) and evaporite (anhydrite, gypsum, halite, and sylvite) minerals showed that most groundwater samples significantly under saturated. The earth's alkaline ions rise as a result, and the predominant carbonate mineral dissolves in the groundwater. In groundwater, evaporite mineral dissolution predominates. The Gibbs diagram illustrates how new groundwater recharges from fresh surface water, with the bulk of the samples being plotted in the rock dominance field. Since some of the samples included saline groundwater, it is likely that the groundwater has been contaminated by dissolved evaporites, older deep water, and/or saline surface water along the flow route. The predominant mineral dissolution found in the groundwater

samples was carbonate and evaporite. The results demonstrated that surface water sources with  $\text{Ca-Mg-HCO}_3$  and  $\text{Na-HCO}_3\text{-Cl}$  water types, as well as the presence of bicarbonate ions in the groundwater, are the primary sources of groundwater recharge. The main controlling factors influencing groundwater salinity are  $\text{SO}_4^{2-}$  ions, earth alkaline elements, and alkali. Due to the ideal limits of TDS,  $\text{Mg}^+$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{SO}_4^{2-}$ , and  $\text{Cl}^-$ , the majority of samples fall within the appropriate range. Residential use of groundwater should be avoided in areas when EC, total hardness,  $\text{Ca}^+$ , and  $\text{NO}_3^-$  slightly exceed recommended standards. It was discovered that the greatest number of water samples was ideal for cultivation.

### 6. REFERENCES

1. N. S. Rao, A. Dinakar, M. Sravanthi, B. K. Kumari, (2021) Geochemical characteristics and quality of groundwater evaluation for drinking, irrigation, and industrial purposes from a part of hard rock aquifer of South India, *Environmental Science and Pollution Research*, **28**: 31941-31961.
2. G. Chandnani, P. Gandhi, D. Kanpariyaa, D. Parikh, M. Shah, (2022) A comprehensive analysis of contaminated groundwater: Special emphasis on nature-ecosystem and socio-economic impacts, *Groundwater for Sustainable Development*, **19**: 100813.
3. G. Ravindiran, S. Rajamanickam, S. Sivarethnamohan, B. Karupaiyasathaiah, G. Ravindran, S. K. Muniasamy, G. Hayder, (2023) A review of the status, effects, prevention, and remediation of groundwater contamination for sustainable environment, *Water*, **15(20)**: 3662.
4. K. Sharma, S. Rajan, S. K. Nayak, (2024) Water pollution: Primary sources and associated human health hazards with special emphasis on rural areas. In: *Water Resources Management for Rural Development*, Netherlands: Elsevier, p3-14.
5. K. M. Ahmed, (2021) Challenges of sustainable groundwater development and management in Bangladesh: Vision 2050. In: *Global Groundwater*, Netherlands: Elsevier. p425-438.
6. R. Panhwar, A. Abro, A. R. Kandhro, N. Khaskheli Jalbani, K. A. Gishkori, S. Qaisar, (2022) Global water mapping, requirements, and concerns over water quality shortages. In: *Water Quality- New Perspectives*, London: IntechOpen.

7. R. Sabale, B. Venkatesh, M. Jose, (2023) Sustainable water resource management through conjunctive use of groundwater and surface water: A review. *Innovative Infrastructure Solutions*, **8(1)**: 17.
8. C. Rout., R. Singh (2024), Water resources management in India: Current state and challenges. In: *National Conference on Science Technology Engineering and Management (STEM-2024)*.
9. R. Ramya Priya, L. Elango, (2022) Geogenic and anthropogenic impacts on the water quality of cauvery river. In: *Riverine Systems: Understanding the Hydrological, Hydrosocial and Hydro-Heritage Dynamics*, Cham: Springer International Publishing, p149-170.
10. L. Kumar, R. Kumari, A. Kumar, I. A. Tunio, C. Sassanelli, (2023) Water quality assessment and monitoring in Pakistan: A comprehensive review, *Sustainability*, **15(7)**: 6246.
11. N. S. Rao, R. Das, H. K. Sahoo, S. Gugulothu, (2024) Hydrochemical characterization and water quality perspectives for groundwater management for urban development, *Groundwater for Sustainable Development*, **24**: 101071.
12. R. Sathre, S. M. Antharam, M. Catena, (2022) Water security in South Asian cities: A review of challenges and opportunities, *CivilEng*, **3(4)**: 873-894.
13. N. Bassi, T. Riefky, Y. Kabir, A. Bhaduri, G. Schmidt, E. Khanduja, M. K. Martawidjaja, (2023) *Improving Access to Safely Managed Drinking Water Services: Recommendations for G20 Emerging Economies*, [T20 Policy Brief].
14. M. Kumar, R. Kumar, (2024) Water management: Issues and challenges. In: *Advances in Water Management Under Climate Change*, United States: CRC Press, p17-34.
15. N. Shukla, S. Gupta, S. Rai, (2023) Potential impacts of climatic changes and human activity on water quality. In: *Environmental Processes and Management: Tools and Practices for Groundwater*, Cham: Springer International Publishing, p103-111.
16. M. Shaikh, F. Birajdar, (2024) Groundwater and ecosystems: Understanding the critical interplay for sustainability and conservation, *EPRA International Journal of Multidisciplinary Research (IJMR)*, **10(3)**: 181-186.
17. B. Panneerselvam, N. Ravichandran, S. P. Kaliyappan, S. Karuppannan, B. Bidorn, (2023) Quality and health risk assessment of groundwater for drinking and irrigation purpose in semi-arid region of India using entropy water quality and statistical techniques, *Water*, **15(3)**: 601.
18. S. Bhatt, A. P. Mishra, N. Chandra, H. S. Sahu, K. Chaurasia, C. B. Pande, J. Hunt, (2024) Characterizing seasonal, environmental and human-induced factors influencing the dynamics of Rispana River's water quality: Implications for sustainable river management, *Results in Engineering*, **22**: 102007.
19. S. Ahmed, M. Mulhim, F. Qureshi, N. Akhtar, S. Lagudu, (2022) Reckoning groundwater quality and hydrogeochemical processes for drinking and irrigation purposes under the influence of anthropogenic activities, North India, *Pollutants*, **2(4)**: 486-509.
20. A. Nayak, G. Matta, D. P. Uniyal, (2023) Hydrochemical characterization of groundwater quality using chemometric analysis and water quality indices in the foothills of Himalayas, *Environment, Development and Sustainability*, **25(12)**: 14229-14260.
21. A. Mukherjee, P. Coomar, S. Sarkar, K. H. Johannesson, A. E. Fryar, M. E. Schreiber, A. Vengosh, (2024) Arsenic and other geogenic contaminants in global groundwater, *Nature Reviews Earth and Environment*, **5**: 312-328.
22. S. Pradhan, S. Chand, P. R. Chand, S. Rout, K. Naik, (2023) Emerging groundwater contaminants: A comprehensive review on their health hazards and remediation technologies, *Groundwater for Sustainable Development*, **20**: 100868.
23. M. Gouasmia, F. Dhahri, H. Azaiez, M. K. Zidi, M. Soussi, (2023) Hydrogeological, geophysical, geochemical and statistical integrated techniques to assess a multilayered groundwater aquifer system in an arid region: The case of the Sbeitla aquifers in Central Tunisia, *Environmental Earth Sciences*, **82(13)**: 320.
24. Y. Hamed, R. Hadji, R. Ahmadi, Y. Ayadi, K. Shuhab, A. Pulido-Bosch, (2024) Hydrogeological investigation of karst aquifers using an integrated geomorphological, geochemical, GIS, and remote sensing techniques (Southern Mediterranean Basin-Tunisia), *Environment, Development and Sustainability*, **26(3)**: 6943-6975.
25. S. P. Rai, K. V. Akpataku, J. Noble, A. Patel, S. K. Joshi, (2023) Hydrochemical evolution of groundwater in northwestern part of the Indo-Gangetic Basin, India: A geochemical and isotopic approach, *Geoscience Frontiers*, **14(6)**: 101676.
26. V. Sivakumar, S. M. Chidambaram, S. Velusamy, R. Rathinavel, D. K. Shanmugasundaram, P. Sundararaj, K. Balu, (2023) An integrated approach for an impact assessment of the tank water and groundwater quality in Coimbatore region of South India: Implication from anthropogenic activities, *Environmental Monitoring and Assessment*, **195(1)**: 88.
27. S. Mishra, M. S. Chauhan, S. Sundaramurthy, (2024) Assessing groundwater quality dynamics in Madhya Pradesh: Chemical contaminants and their temporal patterns, *Environmental Research*, **252**: 118887.
28. N. Maharjan, S. P. Kayastha, C. Bhuiyan, (2024) Hydrogeochemical and microbial analysis of springs in Raghuganga rural municipality of Gandaki Province, Nepal. *Environmental Earth Sciences*, **83(1)**: 13.
29. R. Kausher, R. Singh, A. K. Sinha, S. N. Sathy, S. Kumar, S. Pandey, A. Mohamed, (2023) Assessing impacts of mining-induced land use changes on groundwater and surface water quality using isotopic and hydrogeochemical signatures, *Sustainability*, **15(14)**: 11041.
30. P. Padhi, N. Bora, P. Sohtun, M. Athparia, M. Kumar, R. Katak, P. K. Sarangi, (2024) Remediation of mine overburden and contaminated water with activated biochar derived from low-value biowaste. *Journal of the Taiwan Institute of Chemical Engineers*, **159**: 105472.
31. M. Shehata, S. M. Zaid, S. T. Al-Goul, A. Shami, A. K. M. Al Syaad, A. E. Ahmed,... & S. F. Abu Qamar, (2024) Integrated management of groundwater quantity, physicochemical properties, and microbial quality in West Nile delta using a new MATLAB code and geographic information system mapping, *Scientific Reports*, **14(1)**: 7762.
32. S. Ahmad, R. Umar, I. Ahmad, (2024) Assessment of groundwater quality using Entropy-Weighted Quality Index (EWQI) and multivariate statistical techniques in Central Ganga plain, India, *Environment, Development and Sustainability*, **26(1)**: 1615-1643.
33. P. S. Ganvir, R. Guhey, (2023) An implication of enhanced rock weathering on the groundwater quality: A case study from Wardha Valley Coalfields, Central India. In: *Weathering and Erosion Processes in the Natural Environment*, United States: John Wiley and Sons, p215-242.

34. T. Subramani, P. Gopinathan, V. Satheeskumar, D. Karunanidhi. (2024) Role of geochemistry of rocks and soils for groundwater quality in the Chithar River Basin, South India-an approach using enrichment factor (EF) of trace elements and chemical index of alteration (CIA). In: *Modern River Science for Watershed Management: GIS and Hydrogeological Application*, Cham: Springer Nature Switzerland, p381-402
35. B. K. Nagaraja Rao, S. T. Rajurkar, G. Ramlingaswamy, B. R. Babu, (1987) Stratigraphy, structure and evolution of the Cuddapah basin, *Memoir Geological Society of India*, **6**: 33-87.
36. CGWB, (2016) *Ground Water Year Book of India 2014-15, CGWB, Ministry of WR, RD and GR, Govt of India*. Available from: <https://cgwb.gov.in/ground> [Last accessed on 2024 Nov 02].
37. A. Asmoay, W. A. Mabrouk, (2024) Appraisal of rock-water interaction and frailty of groundwater to corrosion and salinization, northwestern Gulf of Suez, Egypt, *Journal of Umm Al-Qura University for Applied Sciences*, **10(1)**: 39-50.
38. H. T., Abadi, T., Alemayehu, B. A. Berhe, (2024) Hydrogeochemical characterization of groundwater in mountainous catchment and its suitability for drinking purposes in Irob, Tigray, Northern Ethiopia, *Water Practice and Technology*, **19**: 1495.
39. B. Hamma, A. Alodah, F. Bouaicha, M. F. Bekkouche, A. Barkat, E. E. Hussein, (2024) Hydrochemical assessment of groundwater using multivariate statistical methods and water quality indices (WQIs), *Applied Water Science*, **14(2)**: 33.
40. M. A. Sulaiman, M. M. Zafar, R. Prabhakar, R. Kumar, R. K. Sinha, A. Kumari, (2023) A multivariate statistical approach to evaluate the hydro-geochemistry of groundwater quality in the middle Ganga river basin, Patna, India, *Acta Geophysica*, **72**: 1913-1926.
41. A. M. Piper. (1944) A graphical interpretation of water analysis, *Eos, Transactions American Geophysical Union*, **25**: 914-928.
42. C. Arulrajan, M. C. Wu, S. Porada, J. E. Dykstra, C. H. Hou, A. Van der Wal, (2024) Mitigation of mineral scaling in membrane capacitive deionization-Understanding the role of pH changes and carbonates, *Journal of Water Process Engineering*, **60**: 105094.
43. A. Mohamed, F. Asmoay, A. Alshehri, A. Abdelrady, A. Othman, (2022) Hydro-geochemical applications and multivariate analysis to assess the water-rock interaction in arid environments, *Applied Sciences*, **12(13)**: 6340.
44. P. Singh, M. S. Rishi, L. Kaur, (2023) Hydrogeochemical characterization of groundwater under natural and anthropogenically influenced areas located in Upper Ghaggar River Basin, India, *Environment, Development and Sustainability*, **26**: 1-23.
45. G. K. Brhane, H. S. Mekonen, (2024) Identifying controlling factors for the evolution of shallow groundwater chemistry of Ellala catchment, northern Ethiopia, *Sustainable Water Resources Management*, **10(2)**: 1-20.
46. World Health Organization, (2017) *Safely Managed Drinking Water: Thematic Report on Drinking Water*, Switzerland: World Health Organization.
47. H. Rasouli, K. Belhassan, A. Vaseashta, (2024) Hydrogeological investigations of Paghman valleys in Kabul, Afghanistan, *Advances in Geological and Geotechnical Engineering Research*, **6(1)**: 1-20.
48. M. A. Gharaghani, A. Mohammadpour, M. Keshtkar, A. Azhdarpoor, R. Khaksefidi, (2024) Quality evaluation and health risk assessment of drinking water in Minab County: Hydrochemical analysis and artificial neural network modeling, *Environmental Science: Water Research and Technology*, **10(1)**: 250-262.
49. M. Kneuver, D. Mele, R. Sulpizio, (2023) Mineralization and Skarn formation associated with alkaline magma chambers emplaced in a limestone basement: A review, *Minerals*, **13(9)**: 1184.
50. A. Ravindran, A. Promilton, J. V. Kingston, S. R. Abishek, S. K. Aswin, R. Abinaya, R. S. Priya, (2024) Geophysical and Geochemical Studies on Sinking Stream Occurrence in Ayankulam Village of South Tamilnadu, India. In: *Proceedings of the Indian National Science Academy*, p1-15.
51. D. Maji, S. Dutta, (2024) Strategies to safeguard drinking water from hazardous chemical contaminants. In: *Development in Wastewater Treatment Research and Processes*, Netherlands: Elsevier. p419-441.
52. E. Shaji, K. V. Sarath, M. Santosh, P. K. Krishnaprasad, B. K. Arya, M. S. Babu, (2024) Fluoride contamination in groundwater: A global review of the status, processes, challenges, and remedial measures, *Geoscience Frontiers*, **15(2)**: 101734.
53. S. Gupta, N. Mishra, A. Kumar, A. K. Yadav, (2024) Fluoride pollution in subsurface water: Challenges and opportunities. In: *Advanced Treatment Technologies for Fluoride Removal in Water: Water Purification*, Germany: Springer Nature, p19-39.
54. C. D. Aju, A. L. Achu, M. Maharooof, M. C. Raicy, R. Reghunath, G. Gopinath, (2024) Emerging nitrate contamination in groundwater: Changing phase in a fast-growing state of India, *Chemosphere*, **357**: 141964.
55. M. Riaz, M. Ahmad, N. M. Mukhtar, N. Nawaz, (2024) Nitrate contamination of soil and water: Implications for ecosystem functions and human health. In: *Inorganic Contaminants and Radionuclides*, Netherlands: Elsevier, p351-373.

**\*Bibliographical Sketch**

Dr. C. Krupavathi completed Ph.D. as DST-Inspire fellow from Department of science and technology, New Delhi. She completed M.Sc., from Yogi Vemana University, Kadapa. She has received DST-Inspire fellow award in the year of 2018. Her research interests are in the areas of Hydrogeochemistry, Water quality, Contamination of Water, Remediation methods. Her contribution has been recognized in many peer-reviewed journals and book chapters.



Dr. S. Srinivasa Gowd obtained Ph.D. Degree from Sri Venkateswara, Tirupati. He has many publications in the peer reviewed national and international Journals. He is currently working as Associate Professor in the Department of Geology, Yogi Vemana University, Kadapa. He completed more than 16 years of teaching experience in Geology. He has been awarded with CSIR-Research Associate, CSIR-Sr. Research Associate (Scientists' Pool Scheme), and DST-Young Scientist fellowships at National Geophysical Research Institute (NGRI), Hyderabad and Sri Venkateswara University, and Tirupati. His research interests are in the areas of Environmental Geochemistry, Hydrogeochemistry, Remote Sensing and GIS and he has supervised many M.Sc. students' projects.



Dr. P. Ravi Kumar (DST-Inspire Fellow) obtained from Ph.D. Degree from Yogi Vemana University, Kadapa. He completed M.Sc., from Yogi Vemana University, Kadapa. He has many publications in peer-reviewed national and international Journals. He has received the DST-Inspire Fellow Award in the year 2019. His research interests are in the areas of Hydrogeology, Hydrogeochemistry, Remote sensing and GIS, Groundwater Pollution, and Remediation methods.



G. Nagalakshmi Devi completed M.Sc., in 2021 from Yogi Vemana University, Kadapa. She is working as Project Fellow in the RUSA project. Her research interests are in the areas of Hydrogeochemistry, Water quality, Contamination of Water, and Remediation methods.



U. Lahari completed M.Sc., in 2022 from Acharya Nagarjuna University, Guntur. She is working as Research Scholar in the Department of Geology, Yogi Vemana University, Kadapa. Her research interests are in the areas of Hydrogeochemistry, Water quality, Contamination of Water, and Remediation methods.